













# Presentation, care, and outcomes of patients with NSTEMI according to World Bank country income classification: the ACVC-EAPCI EORP NSTEMI Registry of the European Society of Cardiology

Ramesh Nadarajah <sup>1,2,3,\*</sup>, Peter Ludman <sup>4</sup>, Cécile Laroche<sup>5</sup>,  
Yolande Appelman <sup>6</sup>, Salvatore Brugaletta <sup>7,8</sup>, Andrzej Budaj <sup>9</sup>,  
Hector Bueno <sup>10,11,12</sup>, Kurt Huber <sup>13,14</sup>, Vijay Kunadian <sup>15,16</sup>,  
Sergio Leonardi <sup>17,18</sup>, Maddalena Lettino <sup>19</sup>, Dejan Milasinovic <sup>20</sup>  
and Chris P Gale <sup>1,2,3</sup>, on behalf of the NSTEMI investigator group<sup>†</sup>

<sup>1</sup>Leeds Institute for Cardiovascular and Metabolic Medicine, University of Leeds, LS2 3AA, Leeds, UK; <sup>2</sup>Leeds Institute of Data Analytics, University of Leeds, LS2 9NL, Leeds, UK; <sup>3</sup>Department of Cardiology, Leeds Teaching Hospitals NHS Trust, Leeds, UK; <sup>4</sup>Institute of Cardiovascular Sciences, University of Birmingham, Birmingham, UK; <sup>5</sup>EURObservational Research Programme, European Society of Cardiology, European Heart House, Route des Colles, Sophia Antipolis, 2035, France; <sup>6</sup>Department of Cardiology, Amsterdam UMC-Vrije Universiteit, Amsterdam Cardiovascular Sciences, Amsterdam, The Netherlands; <sup>7</sup>Hospital Clinic de Barcelona, Barcelona, Spain; <sup>8</sup>Institut d'Investigacions Biomèdiques August Pi i Sunyer, Barcelona, Spain; <sup>9</sup>Department of Cardiology, Center of Postgraduate Medical Education, Grochowski Hospital, Warsaw, Poland; <sup>10</sup>Cardiology Department, Hospital Universitario 12 de Octubre and Instituto de Investigación Sanitaria Hospital 12 de Octubre (imas12), Madrid, Spain; <sup>11</sup>Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain; <sup>12</sup>Facultad de Medicina, Universidad Complutense de Madrid, Madrid, Spain; <sup>13</sup>3rd Medical Department, Cardiology and Intensive Care Medicine, Clinic Ottakring (Wilhelminenhospital), Vienna, Austria; <sup>14</sup>Sigmund Freud University, Medical Faculty, Vienna, Austria; <sup>15</sup>Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK; <sup>16</sup>Cardiothoracic Centre, Freeman Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK; <sup>17</sup>University of Pavia, Pavia, Italy; <sup>18</sup>Fondazione IRCCS Policlinico S.Matteo, Pavia, Italy; <sup>19</sup>Cardio-Thoracic and Vascular Department, San Gerardo Hospital, ASST-Monza, Monza, Italy; and <sup>20</sup>Department of Cardiology, University Clinical Center of Serbia and Faculty of Medicine, University of Belgrade, Belgrade, Serbia

Received 19 January 2023; revised 23 January 2023; accepted 1 February 2023; online publish-ahead-of-print 3 February 2023

## Background

The majority of NSTEMI burden resides outside high-income countries (HICs). We describe presentation, care, and outcomes of NSTEMI by country income classification.

## Methods and results

Prospective cohort study including 2947 patients with NSTEMI from 287 centres in 59 countries, stratified by World Bank country income classification. Quality of care was evaluated based on 12 guideline-recommended care interventions. The all-or-none scoring composite performance measure was used to define receipt of optimal care. Outcomes included in-hospital acute heart failure, stroke/transient ischaemic attack, and death, and 30-day mortality. Patients admitted with NSTEMI in low to lower-middle-income countries (LLMICs), compared with patients in HICs, were younger, more commonly diabetic, and current smokers, but with a lower burden of other comorbidities, and 76.7% met very high risk criteria for an immediate invasive strategy. Invasive coronary angiography use increased with ascending income classification (LLMICs, 79.2%; upper middle income countries [UMICs], 83.7%; HICs, 91.0%), but overall care quality did not ( $\geq 80\%$  of eligible interventions achieved: LLMICs, 64.8%; UMICs 69.6%; HICs 55.1%). Rates of acute heart failure (LLMICs, 21.3%; UMICs, 12.1%; HICs, 6.8%;  $P < 0.001$ ), stroke/transient ischaemic attack (LLMICs: 2.5%; UMICs: 1.5%; HICs: 0.9%;  $P = 0.04$ ), in-hospital mortality (LLMICs, 3.6%; UMICs: 2.8%; HICs: 1.0%;  $P < 0.001$ ) and 30-day mortality (LLMICs, 4.9%; UMICs, 3.9%; HICs, 1.5%;  $P < 0.001$ ) exhibited an inverse economic gradient.


## Conclusion

Patients with NSTEMI in LLMICs present with fewer comorbidities but a more advanced stage of acute disease, and have worse outcomes compared with HICs. A cardiovascular health narrative is needed to address this inequity across economic boundaries.

<sup>†</sup> Investigator group list at end of manuscript and in separate document.

\* Corresponding author. Tel: +44 (0) 113 343 3241, Email: [r.nadarajah@leeds.ac.uk](mailto:r.nadarajah@leeds.ac.uk)





**Figure 1** STROBE diagram for the EORP NSTEMI registry analytical cohort.

## Statistics

Categorical data were described using numbers and percentages for non-missing data, and continuous data were described using medians, interquartile range (IQR), means, and standard deviations (SD). Comparison between groups was performed using the Kruskal–Wallis test or using Pearson's  $\chi^2$  test or Fisher's exact test if any count was  $<5$ .

To explore for differences in NSTEMI care quality by country wealth, we stratified the cohort into low (LICs), lower-middle (LMICs), upper-middle (UMICs), and high-income countries (HICs) according to the World Bank classification of the 2021 fiscal year.<sup>15</sup> Given the numbers of countries in the study, the LICs and LMICs were combined to form a single larger group of low to lower-middle-income countries (LLMICs). All analyses were conducted using SAS version 9.4, with statistical significance determined at 5%. Findings are reported in accordance with the Strengthening the reporting of observational studies in epidemiology (STROBE) statement.<sup>24</sup>

## Ethics

Favourable ethics approval for the study was granted at the local and/or regional ethics committees or Institutional Review Boards in each country, according to local regulations. National coordinators, in conjunction with local centres, or participating centres managed the approvals of national or regional ethics committees or Institutional Review Boards, according to local regulations.

## Results

### Study population

From the recruited cohort of 3620 participants, 2947 formed the analytical cohort from 59 countries, across four continents and 287 centres (Figure 1, Supplementary material online, Table S1). Most participating centres were teaching research hospitals (45.0%) followed by general hospitals (41.1%) and 58.2% of hospitals had cardiac surgery onsite (Supplementary material online, Table S2). The majority of patients (1761, 59.8%) and participating centres (192, 66.9%) were from HICs. The median (IQR) age and proportion of women was 66 (57–75) years and 30.8%, with NSTEMI patients in HICs being older and more commonly men and Caucasian (Table 1).

Overall, the comorbidity burden of the cohort was high (Table 1). Each of hypertension, hypercholesterolaemia, or ever smoking was prevalent in over half of the patients at time of diagnosis, more than a third had diabetes mellitus and over a quarter had a previous myocardial infarction. Notably, the patients admitted with NSTEMI in HICs were older than those from LLMICs, and had a higher burden of comorbidity, with a three-fold increased prevalence of atrial fibrillation, peripheral vascular disease, prior percutaneous coronary intervention or coronary artery bypass graft surgery and a two-fold increased prevalence of chronic kidney disease and treatment with dialysis. In HIC countries, patients with NSTEMI also had higher rates of heart failure, COPD, hypercholesterolaemia, and cancer, but lower rates of diabetes.

### Presenting characteristics

Patients admitted in LLMICs, compared with patients in UMICs or HICs, had a longer delay to presentation at hospital following their symptom onset and showed signs of more severe disease at time of their presentation (Table 2). That is, the proportion of patients presenting in Killip Class III or IV was three-fold higher in LLMICs than HICs and ongoing chest pain, life-threatening arrhythmias (or cardiac arrest), and haemodynamic instability were at least twice as common. More than three-quarters of patients presenting with NSTEMI in LLMICs met very high risk criteria for an invasive coronary strategy, whereas only a third of patients did so in HICs (Table 2).

### Quality of care

There was geographic variation in the attainment of the 12 measured quality indicators for NSTEMI care (Figure 2). Whilst all patients were recorded as having received an ECG at baseline, only 45% and 46% received aspirin pre-hospitalization and referral for cardiac rehabilitation, respectively. Across the 12 quality indicators 98.3% of patients received at least 40% of the assessed interventions (60.3% of patients received  $\geq 80\%$  of interventions, 38.0% received  $\geq 40\%$  to  $<80\%$  of eligible interventions). By care grouping, 80.2% received  $\geq 80\%$  of eligible interventions related to investigation and invasive coronary strategies, but only 31.2% for lifestyle care interventions.

**Table 1** Baseline characteristics of NSTEMI patients in analytical cohort by country income classification

	Total (n = 2947)	LLMICs (n = 361)	UMICs (n = 825)	HICs (n = 1761)	Missing data, n (%)
Patient characteristics					
Age (years), median (IQR)	66 (57–75)	58 (51–65)	64 (56–73)	68 (59–77)	0
>= 75 years	761 (25.8)	31 (8.6)	177 (21.5)	553 (31.4)	0
>= 80 years	447 (15.2)	16 (4.4)	93 (11.3)	338 (19.2)	0
Women	907 (30.8)	122 (33.8)	259 (31.4)	526 (29.9)	0
Ethnic origin					107 (3.6)
Caucasian	2252 (79.3)	142 (43.2)	553 (69.6)	1557 (90.7)	
Black	41 (1.4)	31 (9.4)	0/795	10 (0.6)	
Asian	278 (9.8)	129 (39.2)	88 (11.1)	61 (3.6)	
Other	269 (9.5)	27 (8.2)	154 (19.4)	88 (5.1)	
Cardiovascular risk factors					
Hypertension	2023 (69.0)	235 (65.3)	597 (72.9)	1191 (67.9)	15 (0.5)
Ever smoker	1591 (56.9)	159 (45.6)	456 (56.6)	976 (59.4)	151 (5.1)
Current smoker	857 (30.7)	128 (36.7)	297 (36.9)	432 (26.3)	
Former smoker	734 (26.3)	31 (8.9)	159 (19.8)	544 (33.1)	
Diabetes Mellitus	1104 (37.7)	174 (48.6)	332 (40.5)	598 (34.2)	19 (0.6)
Hypercholesterolaemia	1427 (51.9)	125 (39.4)	324 (45.3)	978 (57.0)	200 (6.8)
Cardiovascular comorbidities					
Atrial fibrillation	302 (10.3)	13 (3.6)	82 (10.0)	207 (11.8)	17 (0.6)
Heart failure	464 (15.9)	41 (11.5)	163 (20.0)	260 (14.9)	30 (1.0)
Prior myocardial infarction	759 (26.1)	49 (13.7)	277 (34.5)	433 (24.8)	41 (1.4)
Prior PCI	649 (22.1)	31 (8.6)	209 (25.5)	409 (23.3)	14 (0.5)
Prior CABG surgery	217 (7.4)	9 (2.5)	79 (9.6)	129 (7.3)	5 (0.2)
Peripheral vascular disease	243 (8.6)	12 (3.6)	51 (6.6)	180 (10.4)	115 (3.9)
Stroke	204 (6.9)	19 (5.3)	55 (6.7)	130 (7.4)	6 (0.2)
Relevant comorbidities					
Active malignancy	82 (2.8)	8 (2.2)	9 (1.1)	65 (3.7)	37 (1.3)
COPD	234 (8.0)	27 (7.5)	50 (6.1)	157 (9.0)	30 (1.0)
Chronic kidney disease	367 (12.5)	23 (6.4)	108 (13.2)	236 (13.5)	16 (0.5)

CABG, coronary artery bypass grafting; COPD, chronic obstructive pulmonary disease; HICs, high-income countries; LLMICs, low to lower-middle-income countries; PCI, percutaneous coronary intervention; UMICs, upper-middle-income countries.

Across the spectrum of country income classification 100% of patients received an ECG at baseline, and  $\geq 90\%$  each of aspirin, P2Y12 inhibitors and statins at discharge (Figure 2). In contrast, pre-hospitalisation use of aspirin varied substantially (LLMICs, 39.4%; UMICs, 55.6%; HICs, 41.4%). The use of invasive coronary angiography increased with increasing country economic prosperity (LLMICs, 79.2%; UMICs, 83.7%; HICs, 91.0%). A greater proportion of NSTEMI patients in HICs received referral for cardiac rehabilitation than their counterparts in LLMICs (52.0% vs. 39.9%), but a smaller proportion received dietary advice (79.5% vs. 95.0%).

Overall, attainment of the measured quality indicators was higher in LLMICs and UMICs than HICs ( $\geq 80\%$  of eligible interventions achieved: LLMICs, 64.8%; UMICs 69.6%; HICs 55.1%). Whilst the proportion of patients who received  $\geq 80\%$  of eligible interventions related to investigation and invasive coronary strategies was high and increased with ascending income classification (LLMICs, 74.8%; UMICs, 80.1%; HICs, 81.3%), the proportion of patients who received  $\geq 80\%$  of eligible lifestyle care interventions (LLMICs, 33.5%; UMICs, 30.8%; HICs, 30.9%) was uniformly low (Figure 3).

## In-hospital outcomes and 30-day mortality

Adverse outcomes for patients admitted for NSTEMI were more frequent in less economically developed countries, in spite of their younger age at presentation (Figure 4). The rate of acute heart failure (LLMICs, 21.3%; UMICs, 12.1%; HICs, 6.8%;  $P < 0.001$ ), cardiogenic shock (LLMICs, 9.1%; UMICs, 3.2%; HICs, 2.1%;  $P < 0.001$ ), and stroke/transient ischaemic attack (LLMICs: 2.5%; UMICs: 1.5%; HICs: 0.9%;  $P = 0.04$ ) increased down the income classification economic gradient. Crude in-hospital and 30-day mortality were low (1.8% and 2.5%, respectively) but increased with declining country wealth (in-hospital mortality: LLMICs, 3.6%; UMICs: 2.8%; HICs: 1.0%;  $P < 0.001$ ; 30 day mortality: LLMICs, 4.9%; UMICs, 3.9%; HICs, 1.5%;  $P < 0.001$ ).

## Discussion

This prospective international study of 2947 patients with NSTEMI presenting to 1 of 247 recruiting centres from 59 ESC member countries and affiliates provides novel and high resolution insights

**Table 2 Presenting characteristics of NSTEMI patients in analytical cohort by country income classification**

	Total (n = 2947)	LLMICs (n = 361)	UMICs (n = 825)	HICs (n = 1761)	Missing data, n (%)
Clinical presentation					
Symptom onset to hospital admission, median (IQR)	6.3 (2.4–26.6)	8.0 (4.0–44.5)	7.5 (2.3–29.8)	5.6 (2.1–23.7)	11 (0.4)
Heart rate (bpm), median (IQR)	77 (67–90)	86 (76–98)	80 (70–90)	75 (65–86)	7 (0.2)
Systolic Blood pressure (mmHg), median (IQR)	139 (120–155)	130 (110–140)	136 (120–151)	140 (124–158)	7 (0.2)
Cardiac arrest	41 (1.4)	8 (2.2)	14 (1.7)	19 (1.1)	12 (0.4)
Killip class					12 (0.4)
Class I	2351 (80.1)	247 (68.4)	653 (79.2)	1451 (82.9)	
Class II	400 (13.6)	65 (18.0)	120 (14.6)	215 (12.3)	
Class III	151 (5.1)	40 (11.1)	41 (5.0)	70 (4.0)	
Class IV	33 (1.1)	9 (2.5)	10 (1.2)	14 (0.8)	
Ongoing chest pain	1150 (39.3)	248 (69.9)	382 (46.4)	520 (29.8)	23 (0.8)
Life threatening arrhythmia or cardiac arrest	91 (3.1)	20 (5.5)	31 (3.8)	40 (2.3)	3 (0.1)
Acute heart failure	336 (11.5)	78 (21.7)	127 (15.4)	131 (7.5)	13 (0.4)
Mechanical complication of MI	28 (1.0)	11 (3.1)	9 (1.1)	8 (0.5)	9 (0.3)
Haemodynamic instability	147 (5.0)	27 (7.5)	57 (6.9)	63 (3.6)	7 (0.2)
ST segment deviation indicative of ischaemia	1502 (51.1)	239 (66.2)	514 (62.4)	749 (42.6)	5 (0.2)
Laboratory results					
Peak troponin I/Upper reference, median (IQR)	21 (4–100)	39 (6–100)	12 (2–86)	22 (6–111)	1183 (40.1)
Peak troponin T/Upper reference, median (IQR)	11 (4–32)	6 (3–19)	16 (6–39)	10 (3–30)	1842 (62.5)
Haemoglobin (g/L), median (IQR)	138 (123–150)	125 (112–138)	139 (123–150)	140 (126–150)	69 (2.3)
eGFR CG (ml/min/1.73 m <sup>2</sup> ), median (IQR)	77 (57–93)	76 (55–95)	75 (56–92)	77 (58–93)	51 (1.7)
Risk assessment					
GRACE, median (IQR)	135 (113–161)	136 (119–163)	139 (116–163)	133 (111–160)	116 (3.9)
Very-high risk for invasive strategy criterion met	1358 (46.3)	276 (76.7)	450 (54.6)	632 (36.1)	
In-hospital revascularisation (of those eligible)					
Diagnostic coronary angiography	2578 (87.5)	286 (79.2)	690 (83.7)	1602 (91.0)	1 (0.0)
PCI	1791 (60.9)	225 (62.5)	462 (56.2)	1104 (62.7)	
CABG surgery	328 (11.1)	33 (9.2)	110 (13.4)	185 (10.5)	

CABG, coronary artery bypass grafting; HICs, high-income countries, LLMICs, low to lower-middle-income countries; PCI, percutaneous coronary intervention; UMICs, upper-middle-income countries.


into variation in presentation, care, and early outcomes by World Bank country income classification. Our main finding is that patients admitted with NSTEMI in LLMICs, compared with patients in HICs, are younger, have fewer comorbidities, present with a more advanced manifestation of acute disease, and have worse early clinical outcomes. Furthermore, invasive coronary angiography was used less often for patients with NSTEMI in LLMICs than HICs. A cardiovascular health narrative is needed to address this inequity across international economic boundaries.

Disparities in NSTEMI care delivery and outcomes by country income classification are insufficiently characterized in the literature. Previous reports have provided comparisons between countries within the same income classification,<sup>9,11,13</sup> or not stratified by economic prosperity even when including a wide representation of countries.<sup>12,14</sup> The Global Burden of Disease Study and ESC cardiovascular disease statistics have demonstrated divergent burden of disability-adjusted life-years as a result of ischaemic heart disease by country wealth,<sup>8,25</sup> but they do not specifically address NSTEMI or provide high resolution patient-level clinical data. The EORP Acute Coronary Syndrome (ACS) STEMI registry and EU-ROASPIRE (European Action on Secondary and Primary Prevention

by Intervention to Reduce Events) registries have delivered important insights on geographical variation in care delivery across ESC member countries, for ST-segment elevation myocardial infarction (STEMI) care and coronary artery disease primary and secondary prevention, respectively.<sup>26,27</sup>

Patients in LLMICs presented with NSTEMI a decade earlier and with fewer comorbidities than their counterparts in HICs. The earlier manifestation of NSTEMI in LLMICs compared with HICs is likely because key risk factors for development and acute presentation of ischaemic heart disease—such as smoking, blood pressure control, and air pollution—are significantly more burdensome.<sup>8</sup> The adoption of western lifestyles has also contributed to an increasing burden of diabetes mellitus, such that 80% people who have diabetes mellitus live in low- and middle-income countries,<sup>28</sup> and in our cohort almost half of the patients presenting with NSTEMI in LLMICs were diabetic. Furthermore low educational levels in low income countries have been found to be particularly correlated with adverse cardiovascular events, and this may be because medical care (e.g. management of hypertension, diabetes, and secondary prevention) is likely to be poorer in people with the lowest levels of education compared with those with higher levels of education in low-income






countries.<sup>29</sup> Comorbidity burden was higher in patients in HICs and this reflects the notion that additional life years constitute an additional risk for acquiring chronic conditions;<sup>30</sup> that is, amongst ESC member countries and affiliates, premature death is more common and life expectancy shorter in low- and middle-income countries and so individuals may not have the chance to accumulate multiple conditions.<sup>8</sup>

Three-quarters of patients with NSTEMI in LLMICs were defined as very high-risk. Time from symptom onset to hospital admission varied substantially in LLMICs, but a quarter of patients had delays in excess of 45 h. Though we do not have data for the reason for delay, a previous systematic review has demonstrated that pre-hospital emergency services are underdeveloped in low- and middle-income countries, and that use of an ambulance can be cost-prohibitive for most patients who are required to pay out of pocket in countries that lack a national insurance programme.<sup>1</sup> Thus many patients with acute myocardial infarction travel to hospital by public transport and private cars,<sup>31,32</sup> which can be particularly disadvantageous in rural areas.<sup>33</sup> Furthermore, lack of knowledge of symptoms consistent with myocardial ischaemia has been reported as a prominent reason for treatment delay,<sup>34</sup> especially amongst older and illiterate patients.<sup>35</sup>

An invasive strategy was more likely to be employed for NSTEMI patients in more prosperous economies. Observational data has demonstrated that rates of coronary angiography and revascularization during index hospitalization are inversely related to mortality rates in patients admitted for NSTEMI at 28 days, 6 months and one year after comprehensive adjustment.<sup>11,14,36</sup> Accordingly, inter-

ventional technologies have come to dominate cardiology practice for NSTEMI and this creates a financial burden on cardiovascular healthcare delivery.<sup>8</sup> In middle income countries, compared with high income countries, the number of interventional cardiologists is lower by nearly a half and 24/7 catheter laboratory facilities lower by a third.<sup>8</sup> The structural deficit affecting person power and facilities in less wealthy countries inevitably translates into the procedural deficit we observed.

We also observed underuse of cardiac rehabilitation, with only 36 and 40% of NSTEMI patients in UMICs and LLMICs, respectively, being referred. Though cardiac rehabilitation can reduce mortality, morbidity, and hospital re-admissions,<sup>37</sup> it is only available in approximately one-quarter of low-to-middle income countries,<sup>38</sup> and the costs of a typical supervised cardiac rehabilitation model employed in HICs are prohibitive in a low-income country setting.<sup>39</sup> Conversely, provision of guideline-recommended pharmacological treatments on discharge was very high with >95% of patients with NSTEMI in LLMICs receiving dual antiplatelet therapy and statins. This shows that factors such as a skilled workforce,<sup>2</sup> high levels of knowledge transfer from scientific studies to healthcare professionals,<sup>40</sup> and greater uniformity of recommendations from guidelines of national and regional societies,<sup>13</sup> can overcome economic conditions to achieve guideline-recommended care. The ACVC-EAPCI EORP STEMI registry also documented excellent provision of these medications in Europe and the Middle East,<sup>26</sup> demonstrating the widespread acceptance of the importance of these pharmacological therapies for patients with an acute myocardial infarction.




## Limitations

First, because of the observational nature of the data, the study design cannot show a causal relationship, but only an association between country wealth and differences in presentation, care, and outcomes for NSTEMI. Second, the minority of patients were from LLMICs. Third, a greater proportion of participating centres in LLMICs, compared with HICs, were specialty or teaching research hospitals (73% vs. 50%), had PCI capability (89.3% vs. 82.5%), and had cardiac surgery on site (71% vs. 51%) (Supplementary material online, *Table S2*); thus, the selected sites in LLMICs are not truly representative of all centres in those countries, and the disparities in interventional care by income classification may be wider than we report. Fourth, the majority of participants from HICs were Caucasian (91%), whereas only 43% of participants in LLMICs were Caucasian (with 39% Asian and 9% Black), so biological determinants may have influenced the adverse event rates we observed in different income classifications. Fifth, we assessed for variation by country income classification but did not additionally investigate for an effect of income inequality amongst a country's residents, which has been associated with worse cardiovascular outcomes.<sup>41,42</sup> Sixth, the majority of patients included in the study were men, consequently the results cannot be automatically translated to women. Seventh, the representation of low-income countries within the LLMIC stratification was relatively small (Supplementary material online, *Table S1*), so the disparities in care and outcomes for NSTEMI in these countries may be greater

than we observed. Eighth, despite its wide representation, this is not a global study as not all regions are included. Ninth, the number of outcomes was insufficient to allow adjustment of crude rates by baseline differences and confounders.

## Actions to be taken

The findings of this registry imply further steps to be taken into consideration to achieve equity in NSTEMI outcomes across country income classification. All efforts should be made to increase the availability and use of coronary intervention for NSTEMI in LLMICs. An initiative could follow the example of the highly successful 'Stent for Life', which aimed to drive equal access to primary PCI across 19 countries in Europe, and coincided with increasing rates of primary PCI across ESC member states.<sup>26,43</sup> Cardiac rehabilitation could also be made more widely available in LLMICs through the adaptation of models tailored to low resource settings.<sup>39</sup> Policy strategies to reduce the population burden of ischaemic heart disease are required in LLMICs, where there is a trend towards widening disparities in the prevalence of key risk factors—such as smoking, blood pressure, and diabetes mellitus—compared with HICs.<sup>8</sup> The European Association of Preventative Cardiology 'Prevention in your country' initiative provides a comprehensive resource of exemplar initiatives for risk factor modification.<sup>44</sup> Both the ESC and the World Health Organization must continue to advocate to raise the priority accorded to prevention of disease, create health-promoting environments, and



strengthen national capacity, leadership, and governance for disease prevention.<sup>8,45</sup>

## Supplementary material

Supplementary material is available at *European Heart Journal—Quality of Care and Clinical Outcomes* online.

## Acknowledgements

The EORP Oversight Committee, Registry Executive and Steering Committees. The project management (study launch, data collection coordination, data management, and statistical analyses) was conducted by the EURObservational Research Programme (EORP), European Society of Cardiology, Sophia-Antipolis, France: Catherine Bosc and Carole Toulouse as Clinical Project Managers, Sylvie Chanoni and Emanuela Fiorucci as Project Officers; Gagan Chhabra and Sandrine Anglars as Data Managers, Cécile Laroche as Statistical Project Lead and Stateo as Consultant Statistician. Overall activities were coordinated and supervised by Doctor Aldo P. Maggioni (EORP Scientific Coordinator). Special thanks to the Acute Cardiovascular Care Association and the European Association of Percutaneous Cardiovascular Interventions (EAPCI).

## Funding

Since the start of EORP, the following companies have supported the programme: Abbott Vascular Int. (2011–20), Amgen Cardiovascular (2009–18), AstraZeneca (2014–21), Bayer AG (2009–18), Boehringer Ingelheim (2009–19), Boston Scientific (2009–12), the Bristol Myers Squibb and Pfizer Alliance (2011–19), Daiichi Sankyo Europe GmbH (2011–20), the Alliance Daiichi Sankyo Europe GmbH and Eli Lilly and Company (2014–17), Edwards (2016–19), Gedeon

Richter Plc. (2014–16), Menarini Int. Op. (2009–12), MSD-Merck & Co. (2011–14), Novartis Pharma AG (2014–20), ResMed (2014–16), Sanofi (2009–11), Servier (2009–21), Vifor (2019–22).

**Conflict of interest:** H.B., K.H., P.L., R.N., V.K., and Y.A. report no conflict of interest.

C.P.G. reports personal fees from AstraZeneca, Amgen, Bayer, Boehringer-Ingelheim, Daiichi Sankyo, Vifor, Pharma, Menarini, Wondr Medical, Raisio Group, and Oxford University Press. He has received educational and research grants from BMS, Abbott inc., the British Heart Foundation, National Institute of Health Research, Horizon 2020, and from the European Society of Cardiology, outside the submitted work.

S.B. reports personal fees from Boston Scientific, Insight, Lifetech, iVascular, and Abbott Vascular.

D.M. reports personal fees from Abbott, Boston Scientific, Biosensors, and Terumo. He has participated on Data Safety Monitoring Boards or Advisory Boards for Abbott and Boston Scientific.

A.B. reports personal and institutional investigator fees from AstraZeneca, Bristol Myers Squibb/Pfizer, Sanofi Aventis, Novartis, Amgen, Novo Nordisk, and Bayer. He reports personal fees from AstraZeneca, Bristol Myers Squibb/Pfizer, Sanofi Aventis, Novartis, and Bayer. He has participated in Data Safety Monitoring Boards or Advisory Boards for AstraZeneca, Bristol Myers Squibb/Pfizer, Sanofi Aventis, and Bayer. He has received support for attending meetings by Astra Zeneca, Bristol Myers Squibb/Pfizer, Sanofi Aventis, and Bayer.

S.L. reports personal fees from ICON, Chiesi, AstraZeneca, Daiichi Sankyo, Bayer, Bristol Myers Squibb/Pfizer, and Novo Nordisk.

M.L. reports speaker fees from Bristol Myers Squibb/Pfizer, Boehringer-Ingelheim, Novartis and Sanofi, and manuscript writing fees from Edwards Lifesciences. She has also participated as an advisory board member for Sanofi, Bristol Myers Squibb/Pfizer, and Boehringer-Ingelheim.



Y.A. reports a research grant from the Dutch Heart Foundation. Signed COIs available for all authors.

## Data availability

The data underlying this article are subject to an embargo of 12 months from the publication date of the article. Once the embargo expires, the data will be available upon reasonable request to the EORP NSTEMI Executive Committee Chair, with the approval of the EORP Oversight Committee.

## Appendix—ESC EURObservational Research Programme Non-ST-segment elevation myocardial infarction (NSTEMI) Registry Investigator Group

### EORP Oversight Committee

**2016–18:** A. Vahanian, FR (Chair): A. Budaj, PL: N. Dagnes, DE: N. Danchin, FR: V. Delgado, NL: J. Emberson, GB: O. Friberg, SE: C.P. Gale, GB: G. Heyndrickx, BE: B. lung, FR: S. James, SE: A.P. Kappetein, NL: A.P. Maggioni, IT: N. Maniadakis, GR: K.V. Nagy, HU: G. Parati, IT: A-S. Petronio, IT: M. Pietila, FI: E. Prescott, DK: F. Ruschitzka, CH: F. Van de Werf, BE: F. Weidinger, AT: U. Zeymer, DE: **2018–20:** C.P. Gale, GB (Chair): B. Beleslin, RS: A. Budaj, PL: O. Chioncel, RO: N. Dagnes, DE: N. Danchin, FR: J. Emberson, GB: D. Erlinge, SE: M. Glikson, IL: A. Gray, GB: M. Kayikcioglu, TR: A.P. Maggioni, IT: K.V. Nagy, HU: A. Nedoshivin, RU: A-P. Petronio, IT: J.W. Roos-Hesselink, NL: L. Wallentin, SE: U. Zeymer, DE: **2020–22:** B.A. Popescu, RO (Chair): D. Adlam, GB: A.L.P. Caforio, IT: D. Capodanno, IT: M. Dweck, GB: D. Erlinge, SE: M. Glikson, IL: J. Hausleiter, DE: B. lung, FR: M. Kayikcioglu, TR: P. Ludman, GB: L. Lund, SE: A.P. Maggioni, IT: S. Matskeplishvili, RU: B. Meder, DE: K.V. Nagy, HU: A. Nedoshivin, RU: D. Neglia, IT: A.A. Pasquet, BE: J.W. Roos-Hesselink, NL: F.J. Rossello, ES: S.M. Shaheen, EG: A. Torbica, IT.

### Executive Committee

Christopher Peter Gale (Chair), Peter F Ludman (Chair), Maddalena Lettino, Hector Bueno, Kurt Huber, Sergio Leonardi, Andrzej Budaj, Dejan Milasinovic (Serbia), Salvatore Brugaletta, Yolande Appelman, Vijay Kunadian

### National Coordinators

AE: Wael Abdul Rahman Al Mahmeed, AM: Hovhannes Kzhdryan, AR: Carlos Dumont, AT: Alexander Geppert, BA: Nirvana Sabanovic Bajramovic, BD: Fathima Aaysha Cader, BE: Christophe Beauloye, CR: Daniel Quesada, CZ: Ota Hlinomaz, DE: Christoph Liebetau, EE: Toomas Marandi, EG: Khaled Shokry, ES: Héctor Bueno, FR: Jacques Mansourati, GB: Chris P Gale, GE: David Maisuradze, GR: Theodoros Sinanis, ID: Surya Dharma, IL: Katia Orvin, IN: Ajay Sinha, IQ: Hasan Ali Farhan, IR: Parham Sadeghipour, IS: Ingibjorg Gudmundsdottir, IT: Ciro Indolfi, JP: Takeshi Kitai, KG: Erkin Mirrakhimov, LT: Olivija Gustiene, LU: François Philippe, LV: Andrejs Erglis, MD: Mihail Popovici, MK: Oliver Kalpak, MN: Mungunchimeg Dagva, MT: Robert George Xuereb, MY: Wan Azman Wan Ahmad, NL: Yolande Appelman, OM: Sunil K. Nadar, PK: Nadeem Qamar, PL: Marek Gierlotka, PT: Silvia Monteiro, RO: Gheorghe-Andrei Dan, RS: Mina Radovanovic, RU: Elena Vladimirovna Reznik, SA: Mohammed Balghith, SD: Awad Mohamed, SI: Miha Cercek, SK: Denisa Brozmannová, TR: Vedat Aytekin, UG: Prossie Ingabire, UZ: Baxrom Alyavi.

## Investigators

**Argentina:** *La Plata:* L. Cartasegna, V. Nomberto Gomez, C. Beltrano, V. Novas, *Rosario:* C. Dumont, H. Balparda, M. Cardona, C. Cigalini, V. Hoyle, *Rosario:* E. San Dámaso, V. Tapia, **Armenia:** *Yerevan:* K. Poghosyan, S. Mkhitarian, M. Adamyan, S. Boyadjian, *Yerevan:* H. Hayrapetyan, K. Azaryan, M. Tadevosyan, H. Poghosyan, *Yerevan:* H. Kzhdryan, A. Vardanyan, **Austria:** *Vienna:* K. Huber, A. Geppert, A. Ahmed, *Vienna:* F. Weidinger, M. Derntl, M. Hasun, T. Schuh-Eiring, L. Riegler, **Bangladesh:** *Dhaka:* M.M. Haq, F.A. Cader, M.A-M. Dewan, M-E. Fatema, A.S. Hasan, M.M. Islam, F. Khandoker, R. Mayedah, S.U. Nizam, *Dhaka:* M.G. Azam, M.M. Arefin, J. Jahan, **Belgium:** *Aalst:* D. Schelfaut, H. De Raedt, S. Wouters, S. Aerts, H. Batjoens, *Brussels:* C. Beauloye, M. Dechamps, S. Pierard, O. Van Caenegem, F. Sinnaeve, *Edegem:* M.J. Claeys, M. Snepvangers, V. Somers, *Gent:* S. Gevaert, H. Schaubroeck, P. Vervaet, M. Buysse, *Kortrijk:* F. Renders, M. Dumoulein, N. Hiltrop, M. De Coninck, S. Naessens, I. Senesael, *Liege:* E. Hoffer, S. Pourbaix, J. Beckers, C. Dugauquier, S. Jacquet, D. Malmendier, M. Massoz, *Liege:* P. Evrard, L. Collard, *Mons:* P. Brunner, S. Carlier, M. Blockmans, D. Mayne, E. Timiras, *Yvoir:* A. Guédès, F. Demeure, C. Hanet, J. Domange, K. Jourdan, **Bosnia and Herzegovina:** *Sarajevo:* E. Begic, F. Custovic, A. Dozic, E. Hrvat, I. Kurbasic, D. Mackic, A. Subo, *Sarajevo:* A. Durak-Nalbantic, A. Dzubur, D. Rebic, A. Hamzic-Mehmedbasic, A. Redzepovic, A. Djokic-Vejzovic, E. Hodzic, M. Hujdur, E. Musija, Z. Gljiva-Gogic, N. Serdarevic, *Sarajevo:* N. Sabanovic Bajramovic, L. Brigic, M. Halilcevic, M. Cibo, N. Hadžibegic, N. Kukavica, *Sarajevo:* A. Begic, A. Iglica, A. Osmanagic, N. Resic, M. Vucijak Grgurevic, F. Zvizdic, *Zenica:* B. Pojskic, E. Mujaric, H. Selimovic, M. Ejubovic, L. Pojskic, E. Stimjanin, M. Sut, **Colombia:** *Medellin:* P. Sanchez Zapata, C. Gallego Munoz, L.A. Fajardo Andrade, M.P. Tobon Upegui, L.E. Perez, **Costa Rica:** *Heredia:* J. Chavarria, D. Quesada, K. Alvarado, **Croatia:** *Rijeka:* L. Zaputovic, V. Tomulic, D. Gobic, T. Jakljevic, D. Lulic, G. Bacic, L. Bastiancic, **Cyprus:** *Nicosia:* P. Avraamides, C. Eftychiou, N. Eteocleous, A. Ioannou, C. Lambrianidi, M. Drakomath-ioulakis, **Czech Republic:** *Brno:* L. Groch, O. Hlinomaz, M. Rezek, J. Semenka, J. Sitar, M. Beranova, P. Kramarikova, *Ceske Budejovice:* L. Pesl, S. Sindelarova, F. Tousek, **Egypt:** *Alexandria:* H.M. Warda, I. Ghaly, S. Habiba, A. Habib, *Aswan:* M.N. Gergis, H. Bahaa, A. Samir, *Cairo:* H. Salah Eldin Taha, M. Adel, H. Mahrous Algamal, M. Mamdouh, *Cairo:* A. Foad Shaker, K. Shokry, A. Konsoah, A. Magdy Mostafa, A. Ibrahim, *Cairo:* A. Imam, B. Hafez, A. Zahran, *Cairo:* M. Abdelhamid, K. Mahmoud, A. Mostafa, A. Samir, M. Abdrabou, A. Kamal, S. Sallam, *Minya:* A. Ali, K. Maghraby, A. Radi Atta, *Zagazig:* A. Saad, M. Ali, **Estonia:** *Tallinn:* E-M. Lotman, R. Lubi, H. Kaljumäe, *Tallinn:* T. Uuetoa, U. Kiitam, **France:** *Argenteuil:* C. Durier, *Auxerre:* O. Ressenecourt, A. Alaa El Din, A. Guiatni, M. Le Bras, E. Mougenot, *Bayonne:* J-N. Labeque, J-L. Banos, O. Capendeguy, *Brest:* J. Mansourati, A. Fofana, M. Augagneur, L. Bahon, A. Le Pape, *Chambery:* L. Batiats-Moreau, A. Fluttaz, F. Good, F. Prieur, *La Roche-Sur-Yon:* E. Boiffard, A-S. Derien, I. Drapeau, N. Roy, *Lyon:* T. Perret, O. Dubreuil, S. Ranc, S. Rio, *Marseille:* J-L. Bonnet, G. Bonnet, T. Cuisset, P. Deharo, J-P. Mouret, J-C. Spychaj, A. Blondelon, *Pau:* N. Delarche, *Pontoise:* V. Decalf, N. Guillard, A. Hakme, M-P. Roger, *Rennes:* Y. Biron, P. Druelles, C. Loubeyre, A. Luccon, P. Hery, *Saint Denis:* M. Nejari, F. Digne, *Saint Nazaire:* F. Huchet, *Thonon-Les-Bains:* A. Neykova, B. Tzvetkov, M. Larrieti, *Toulon:* G. Quaino, P. Armangau, *Toulouse:* A. Sauguet, L. Bonfils, N. Dumonteil, J. Fajadet, B. Farah, B. Honton, B. Monteil, R. Philippart, D. Tchetche, M. Cottin, F. Petit, A. Piquart, *Vandoeuvre-Les-Nancy:* B. Popovic, J. Varlot, **Georgia:** *Tbilisi:* D. Maisuradze, E. Sagirashvili, *Tbilisi:* Z. Kereselidze, L. Totladze, T. Ginturi, D. Lagvilava, **Germany:** *Bad Nauheim:* C. Hamm, C. Liebetau, M. Haas, C. Hamm, T. Koerschgen, M. Weferling, J-S. Wolter, K. Maier, *Bonn:* G. Nickenig, A. Sedaghat, C. Zachoal, **Greece:** *Athens:* K. Lampropoulos, A. Mpatsoili, A. Sakellaropoulou, K. Tyrovolas, N. Zibounomi, K. Argyropoulos, F. Toulgaridis, *Athens:* A. Kolyviras, G. Tzanis, V. Tzifos, *Athens:* A. Milkas, S. Papaioannou,

K. Kyriazopoulos, *Athens*: V. Pylarinou, I. Kontonassakis, C. Kotakos, G. Kourgiannidis, P. Ntoliou, N. Parzakonis, A. Pipertzi, A. Sakalidis, C.-L. Ververeli, *Chania*: K. Kafkala, T. Sinanis, G. Diakakis, K. Grammatikopoulos, E. Papoutsaki, T. Patialiatos, M. Mamaloukaki, S.T. Papadaki, *Serres*: I.E. Kanellos, A. Antoniou, G. Tsinopoulos, C. Goudis, M. Giannadaki, S. Daios, M. Petridou, P. Skantzis, P. Koukis, F. Dimitriadis, M. Savvidis, *Thessaloniki*: I. Styliadis, V. Sachpekidas, A. Pilalidou, N. Stamatiadis, A. Fotoglidis, A. Karakanas, **Hungary**: *Budapest*: Z. Ruzsa, D. Becker, F. Nowotta, **Iceland**: *Reykjavik*: I. Gudmundsdottir, B. Libungan, F. Bjork Skuladottir, H. Halldorsdottir, **India**: *Bangalore*: R. Shetty, S. Iyengar, C. Bs, S. G, S. Lakshmana, R. S, *Patna*: N. Tripathy, A. Sinha, B. Choudhary, A. Kumar, A. Kumar, R. Raj, R. Sagor Roy, **Indonesia**: *Jakarta*: S. Dharmar, B.B. Siswanto, **Iraq**: *Baghdad*: H. Ali Farhan, I.F. Yaseen, *Baghdad*: M. Al-Zaidi, Z. Dakhil, *Erbil*: S. Amen, B. Rasool, *Najaf*: A. Rajeeb, K. Amber, H. Hameed Ali, *Nasiriyah*: T. Al-Kinani, M. Hashim Almyahi, F. Al-Obaidi, **Islamic Republic of Iran**: *Isfahan*: G. Masoumi, M. Sadeghi, K. Heshmat-Ghahdarjani, H. Roohafza, N. Sarrafzadegan, M. Shafeie, Z. Teimouri-Jervekani, F. Noori, *Tehran*: M. Kyavar, P. Sadeghipour, A. Firouzi, M.J. Alemzadeh-Ansari, B. Ghadroost, R. Golpira, A. Ghorbani, F. Ahangari, *Tehran*: M. Salarifar, Y. Jenab, A. Biria, S. Haghighi, P. Mansouri, S. Yadangi, **Israel**: *Petah-Tikva*: R. Kornowski, K. Orvin, A. Eisen, N. Oginez, R. Vazel, H. Kfir, **Italy**: *Bologna*: G. Di Pasquale, G. Casella, L.S. Cardelli, E. Filippini, S. Zagnoni, *Bolzano*: L. Donazzan, D. Ermacora, *Catanzaro*: C. Indolfi, A. Polimeni, A. Curcio, A. Mongiardo, S. De Rosa, S. Sorrentino, C. Spaccarotella, *Crema*: M. Landolina, M. Marino, M. Cacucci, L. Vailati, *Genoa*: P. Bernabò, *Mon serrato*: R. Montisci, L. Meloni, M.F. Marchetti, M. Biddau, E. Garau, *Naples*: E. Barbato, C. Morisco, T. Strisciungio, G. Canciello, *Nuora*: G. Lorenzoni, G. Casu, P. Merella, *Palermo*: G. Novo, A. D'Agostino, D. Di Lisi, A. Di Palermo, S. Evola, F. Immordino, L. Rossetto, G. Spica, *Pordenone*: D. Pavan, A. Della Mattia, R. Belfiore, U. Grandis, F. Vendrametto, C. Spagnolo, L. Carniel, E. Sonego, *Rome*: C. Gaudio, F. Barillà, F.G. Biccire, N. Bruno, I. Ferrari, V. Paravati, C. Torromeo, *Salerno*: G. Galasso, A. Peluso, C. Prota, I. Radano, R.M. Benventa, D. Ferraioli, *San Bonifacio*: M. Anselmi, G.M. Frigo, *Trieste*: G. Sinagra, M. Merlo, A. Perkan, F. Ramani, A. Altinier, E. Fabris, *Turin*: M. Rinaldi, T. Usmiani, L. Checco, S. Frea, M. Mussida, **Japan**: *Fukuoka*: R. Matsukawa, *Hidaka City*: K. Sugi, *Kobe*: T. Kitai, Y. Furukawa, A. Masumoto, Y. Miyoshi, *Shimotsugagun-Tochigi*: S. Nishino, **Kazakhstan**: *Almaty*: B. Assembekov, B. Amirov, Y. Chernokurova, F. Ibragimova, **Kyrgyzstan**: *Bishkek*: E. Mirrakhimov, A. Ibraimova, T. Murataliev, Z. Radzhapova, E. Sultan Uulu, N. Zhanyshbekova, V. Zventsova, **Latvia**: *Riga*: A. Erglis, L. Bondare, **Lithuania**: *Kaunas*: R. Zaliunas, O. Gustiene, R. Dirsiene, J. Marcinkeviciene, G. Sakalyte, A. Virbickiene, G. Baksyte, L. Bardauskiene, R. Gelmaniene, A. Salkauskaite, G. Ziubryte, *Siuliai*: N. Kupstyte-Kristapone, *Vilnius*: J. Badariene, S. Balciute, L. Kapleriene, M. Lizaitis, J. Marinskiene, A. Navickaite, A. Pilkiene, D. Ramauskaite, R. Serpytis, D. Silinskiene, T. Simbelyte, J. Staigyte, **Luxembourg**: *Luxembourg*: F. Philippe, P. Degrell, E. Camus, **Malaysia**: *Kuala Lumpur*: W.A. Wan Ahmad, Z. Abu Kassim, **Malta**: *Maida*: R.G. Xuereb, L.-L. Buttigieg, W. Camilleri, E. Pillaha, S. Xuereb, **Moldova, Republic Of**: *Chisinau*: M. Popovici, V. Ivanov, A. Plugaru, V. Moscalu, I. Popovici, M. Abras, L. Ciobanu, N. Litvinenco, S. Fuior, C. Dumanski, M. Ivanov, T. Danila, *Chisinau*: L. Grib, S. Filimon, L. Cardaniuc, *Chisinau*: A. Batrinac, M. Tasnic, C. Cozma, V. Revenco, *Chisinau*: G. Sorici, **Mongolia**: *Ulaanbaatar*: M. Dagva, G. Choijiljav, E. Dandar, M.-U. Khurelbataar, B. Tsognemekh, **Netherlands**: *Amsterdam*: J. Appelman, A. Den Hartog, H. Jan Te Kolste, D. Van Den Buijs, *Heerlen*: A. Van'T Hof, T. Pustjens, V. Houben, I. Kasperski, *Nieuwegein*: J. Ten Berg, J. Azzahafi, W. Bor, D. Chan Pin Yin, **Nigeria**: *Lagos*: A. Mbakwem, C. Amadi, O. Kushimo, M. Kilasho, E. Oronsaye, **North Macedonia**: *Ohrid*: N. Bakrakeski, E. Kovacheska Bashuroska, V. Mojsovska, S. Tupare, M. Dejan, J. Jovanoska, D. Razmoski, T. Marinovski, *Skopje*: A. Antovski, Z. Jovanovski, S. Kocho, R. Markovski, V. Ristovski, A.B. Samir, S. Biserka, *Skopje*: O. Kalpak, I. Mitevska Peovska, B. Zafirovska Taleska, H. Pejkov, O. Busljetic, Z. Zimbakov, E. Grueva, I. Bojovski, M. Tutic, *Skopje*: V. Poposka, M. Vavlukis, **Oman**: *Muscat*: A. Al-Riyami, S.K. Nadar, W. Abdelmottaleb, S. Ahmed, M. Shaikh Mujtaba, S. Al-Mashari, H. Al-Riyami, **Pakistan**: *Karachi*: A.H. Laghari, O. Faheem, S. Waqar Ahmed, *Karachi*: N. Qamar, S. Furnaz, K. Kazmi, T. Saghir, A. Aneel, A. Asim, F. Madiha, **Poland**: *Bialystok*: B. Sobkowicz, A. Tycinska, E. Kazimierczyk, A. Szyszkowska, *Katowice*: K. Mizia-Stec, M. Wybraniec, A. Bednarek, K. Glowacki, *Kedzierzyn-Kozle*: J. Prokopczuk, W. Babiniski, A. Blachut, M. Kosiak, A. Kusinska, S. Samborski, J. Stachura, H. Szastok, A. Wester, D. Bartoszewska, *Kielce*: B. Sosnowska-Pasiarska, M. Krzysiek, *Krakow*: J. Legutko, B. Nawrotek, *Lodz*: J.D. Kasprzak, M. Klosinska, K. Wiklo, *Lodz*: M. Kurpesa, T. Rechcinski, U. Cieslik-Guerra, *Opole*: M. Gierlotka, J. Bugajski, P. Feusette, J. Sacha, P. Przybylo, *Warsaw*: P. Krzesinski, R. Ryczek, A. Karasek, A. Kazmierczak-Dziuk, M. Mielniczuk, K. Betkier-Lipinska, *Warsaw*: M. Roik, A. Labyk, M. Krakowian, M. Machowski, M. Paczynska, M. Potepa, P. Pruszczuk, *Warsaw*: A. Budaj, M. Ambroziak, E. Omelanczuk-Wiech, A. Torun, *Warsaw*: G. Opolski, R. Glowczynska, A. Fojt, R. Kowalik, *Warsaw*: Z. Huczek, S. Jedrzejszyk, *Wroclaw*: T. Roleder, K. Brust, *Zabrze*: M. Gasiorek, P. Desperak, M. Hawranek, **Portugal**: *Amadora*: P. Farto-Abreu, M. Santos, S. Baptista, L. Brizida, D. Faria, J. Loureiro, P. Magno, C. Monteiro, M. Nédio, *Barreiro*: J. Tavares, C. Sousa, I. Almeida, S. Almeida, H. Miranda, H. Santos, A.P. Santos, *Coimbra*: L. Goncalves, S. Monteiro, R. Baptista, C. Ferreira, J. Ferreira, F. Goncalves, C. Lourenço, P. Monteiro, *Evora*: B. Picarra, A.R. Santos, R. Azevedo Guerreiro, M. Carias, M. Carrington, J. Pais, M. Paralta de Figueiredo, A.R. Rocha, *Faro*: J. Mimoso, I. De Jesus, R. Fernandes, J. Guedes, T. Mota, *Lisbon*: M. Mendes, J. Ferreira, A. Tralhão, C. Tavares Aguiar, C. Strong, F. Fernandes Da Gama, G. Pais, *Lisbon*: A.T. Timóteo, S. Aguiar Oliveira Rosa, T. Mano, J. Reis, M. Selas, *Loures*: D. Espregueira Mendes, M. Satendra, *Penafiel*: P. Pinto, C. Queirós, I. Oliveira, L. Reis, I. Cruz, R. Fernandes, *Porto*: S. Torres, A. Luz, A. Campinas, R. Costa, A. Frias, M. Oliveira, *Santarem*: V. Martins, B. Castilho, C. Coelho, A.R. Moura, N. Cotrim, *Vila Franca de Xira*: R. Carvalho Dos Santos, P. Custodio, R. Duarte, R. Gomes, F. Matias, C. Mendonca, J. Neiva, C. Rabacal, A. Rocha Almeida, *Vila Nova de Gaia*: D. Caeiro, P. Queiroz, G. Silva, **Romania**: *Arad*: A.L. Pop-Moldovan, D. Darabantiu, S. Mercea, *Bucharest*: G.-A. Dan, A.-R. Dan, M. Dobranici, R.A. Popescu, C. Adam, *Bucharest*: C.-J. Sinescu, C.-L. Andrei, R. Brezeanu, N. Samoila, *Bucharest*: M.M. Baluta, *Cluj-Napoca*: D. Pop, R. Tomoaia, *Craiova*: O. Istratoaie, I. Donoiu, A. Cojocar, O.C. Oprita, A. Rocsoreanu, *Iasi*: M. Grecu, S. Ailoeai, *Oradea*: M. Ioachim Popescu, A. Cozma, E. Emilia Babes, M. Rus, A. Ardelean, R. Larisa, M. Moisi, E. Ban, A. Buzle, G. Filimon, *Targu-Mures*: D. Doboreanu, S. Lupu, A. Mitre, R. Rudzik, I. Sus, D. Opris, C. Somkereki, *Timisoara*: C. Mornos, L. Petrescu, A. Betiu, A. Volcescu, O. Ioan, C. Luca, D. Maximov, S. Mosteoru, L. Pascalau, C. Roman, D. Brie, S. Crisan, C. Erimescu, L. Falnita, D. Gaita, M. Gheorghiu, **Russian Federation**: *Chelyabinsk*: S. Levashov, M. Redkina, *Kaluga*: N. Novitskii, E. Dementiev, A. Baglikov, *Moscow*: D. Zateyshchikov, E. Zubova, A. Rogozhina, *Moscow*: A. Salikov, I. Nikitin, E.V. Reznik, M.S. Komissarova, M. Shebzukhova, K. Shitaya, S. Stolbova, *Moscow*: V. Larina, F. Akhmatova, G. Chuvarayan, *Moscow*: M.N. Arefyev, O.V. Averkov, A.L. Volkova, M.S. Sepkhanyan, V.I. Vecherko, *Moscow*: I. Meray, L. Babaeva, L. Goreva, A. Pisaryuk, P. Potapov, M. Teterina, *Moscow*: F. Ageev, G. Silvestrova, *Moscow*: Y. Fedulaev, T. Pinchuk, *Moscow*: I. Staroverov, D. Kalimullin, T. Sukhinina, N. Zhukova, *Tomsk*: V. Ryabov, E. Kruchinkina, D. Vorobeve, *Voronezh*: I. Shevchenko, V. Budyak, O. Elistratova, E. Fetisova, R. Islamov, E. Ponomareva, **Saudi Arabia**: *Hail*: H. Khalaf, A. Al Shaimaa, W. Kamal, *Jeddah*: J. Alrahimi, A. Elshiekh, *Riyadh*: M. Balghith, A. Ahmed, N. Attia, A.A. Jamiel, **Serbia**: *Belgrade*: T. Potpara, M. Marinkovic, M. Mihajlovic, N. Mujovic, A. Kocijancic, Z. Mijatovic, *Belgrade*: M. Radovanovic, D. Matic, A. Milosevic, L. Savic, I. Subotic, A. Uscumlic, N. Zlatic, J. Antonijevic, O. Vesic, *Kragujevac*: R. Vucic, *Nis*: S. Salinger Martinovic,





5. Hall M, Bebb OJ, Dondo TB, Yan AT, Goodman SG, Bueno H et al. Guideline-indicated treatments and diagnostics, GRACE risk score, and survival for non-ST elevation myocardial infarction. *Eur Heart J* 2018;**39**:3798–3806.
6. Schiele F, Gale CP, Simon T, Fox KA, Bueno H, Lettino M et al. The 2020 ESC-ACVC quality indicators for the management of acute myocardial infarction applied to the FAST-MI registries. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:207–215.
7. Collet J-P, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL et al. 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: the Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2021;**42**:1289–1367.
8. Timmis A, Townsend N, Gale CP, Torbica A, Lettino M, Petersen SE et al. European Society of Cardiology: cardiovascular disease statistics 2020. *Eur Heart J* 2020;**41**:12–85.
9. Edfors R, Jernberg T, Lewinter C, Blöndal M, Eha J, Löveke P et al. Differences in characteristics, treatments and outcomes in patients with non-ST-elevation myocardial infarction: novel insights from four national European continuous real-world registries. *Eur Heart J Qual Care Clin Outcomes* 2022;**8**:429–436.
10. Franken M, Giugliano RP, Goodman SG, Baracoli LM, Godoy LC, Furtado RH et al. Performance of acute coronary syndrome approaches in Brazil: a report from the BRACE (Brazilian Registry in Acute Coronary syndromes). *Eur Heart J Qual Care Clin Outcomes* 2020;**6**:284–292.
11. Chung S-C, Sundström J, Gale CP, James S, Deanfield J, Wallentin L et al. Comparison of hospital variation in acute myocardial infarction care and outcome between Sweden and United Kingdom: population based cohort study using nationwide clinical registries. *BMJ* 2015;**351**.
12. Bueno H, Rossello X, Pocock S, Van de Werf F, Chin CT, Danchin N et al. Regional variations in hospital management and post-discharge mortality in patients with non-ST-segment elevation acute coronary syndrome. *Clinical research in cardiology* 2018;**107**:836–844.
13. McNamara R, Chung S, Jernberg T, Holmes D, Roe M, Timmis A et al. International comparisons of the management of patients with non-ST segment elevation acute myocardial infarction in the United Kingdom, Sweden, and the United States: the MINAP/NICOR, SWEDEHEART/RIKS-HIA, and ACTION Registry-GWTG/INCDR registries. *Int J Cardiol* 2014;**175**:240–247.
14. Bueno H, Rossello X, Pocock SJ, Van de Werf F, Chin CT, Danchin N et al. In-hospital coronary revascularization rates and post-discharge mortality risk in non—ST-segment elevation acute coronary syndrome. *J Am Coll Cardiol* 2019;**74**:1454–1461.
15. The World Bank. *World Bank Country and Lending Groups*. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519> (01 July 2021; date last accessed).
16. Nadarajah R, Ludman P, Appelmann Y, Brugaletta S, Budaj A, Bueno H et al. Cohort profile: the ESC EURObservational Research Programme Non-ST-segment elevation myocardial infarction (NSTEMI) Registry. *Eur Heart J Qual Care Clin Outcomes* **9**:8–15, 2022.
17. Roffi M, Patrono C, Collet J-P, Mueller C, Valgimigli M, Andreotti F et al. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Kardiol Pol* 2015;**73**:1207–1294.
18. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA et al. Fourth universal definition of myocardial infarction (2018). *J Am Coll Cardiol* 2018;**72**:2231–2264.
19. Schiele F, Gale CP, Bonnefoy E, Capuano F, Claeys MJ, Danchin N et al. Quality indicators for acute myocardial infarction: a position paper of the Acute Cardiovascular Care Association. *Eur Heart J Acute Cardiovasc Care* 2017;**6**:34–59.
20. Schiele F, Aktaa S, Rossello X, Ahrens I, Claeys MJ, Collet J-P et al. 2020 Update of the quality indicators for acute myocardial infarction: a position paper of the Association for Acute Cardiovascular Care: the study group for quality indicators from the ACVC and the NSTEMI-ACS guideline group. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:224–233.
21. Hackshaw A, Morris JK, Boniface S, Tang J-L, Milenković D. Low cigarette consumption and risk of coronary heart disease and stroke: meta-analysis of 141 cohort studies in 55 study reports. *BMJ* 2018;**360**.
22. Freeman AM, Morris PB, Barnard N, Esselstyn CB, Ros E, Agatston A et al. Trending cardiovascular nutrition controversies. *J Am Coll Cardiol* 2017;**69**:1172–1187.
23. Mehran R, Rao SV, Bhatt DL, Gibson CM, Caixeta A, Eikelboom J et al. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium. *Circulation* 2011;**123**:2736–2747.
24. Von Elm E, Altman D, Egger M, Pocock S, Gøtzsche P, Vandenbroucke J. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Initiative. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet North Am Ed* 2007;**370**:1453–1457.
25. Vos T, Lim SS, Abbafati C, Abbas KM, Abbasi M, Abbasifard M et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet North Am Ed* 2020;**396**:1204–1222.
26. Zeymer U, Ludman P, Danchin N, Kala P, Laroche C, Sadeghi M et al. Reperfusion therapies and in-hospital outcomes for ST-elevation myocardial infarction in Europe: the ACVC-EAPCI EORP STEMI Registry of the European Society of Cardiology. *Eur Heart J* 2021;**42**:4536–4549.
27. Kotseva K, De Backer G, De Bacquer D, Rydén L, Hoeks A, Grobbee D et al. Lifestyle and impact on cardiovascular risk factor control in coronary patients across 27 countries: Results from the European Society of Cardiology ESC-EORP EUROASPIRE V registry. *Eur J Prev Cardiol* 2019;**26**:824–835.
28. Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB et al. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res Clin Pract* 2022;**183**:109119.
29. Rosengren A, Smyth A, Rangarajan S, Ramasundarahettige C, Bangdiwala SI, AlHabib KF et al. Socioeconomic status and risk of cardiovascular disease in 20 low-income, middle-income, and high-income countries: the Prospective Urban Rural Epidemiologic (PURE) study. *Lancet Glob Health* 2019;**7**:e748–e760.
30. Violan C, Foguet-Boreu Q, Flores-Mateo G, Salisbury C, Blom J, Freitag M et al. Prevalence, determinants and patterns of multimorbidity in primary care: a systematic review of observational studies. *PLoS One* 2014;**9**:e102149.
31. Doddipalli SR, Rajasekar D, Vanajakshamma V, Naik KS. Determinants of total ischemic time in primary percutaneous coronary interventions: a prospective analysis. *Indian Heart J* 2018;**70**:S275–S279.
32. Kim D-Y, Wala Z, Islam S, Islam R, Ahn M. Clinical characteristics and outcomes of ST-segment elevation myocardial infarction in a low income setting in rural Bangladesh. *Int J Cardiol Heart Vasc* 2019;**23**:100376.
33. Gupta R, Yusuf S. Challenges in management and prevention of ischemic heart disease in low socioeconomic status people in LLMICs. *BMC medicine* 2019;**17**:1–11.
34. Venkatesan VCK, Madhavi S, Kuzhanthai P. A study to explore the factors related to treatment seeking delay among adults diagnosed with acute myocardial infarction at KMCH, Coimbatore. *Indian Heart J* 2018;**70**:793–801.
35. Sriha Belgith A, Beltaief K, Msolli MA, Bouida W, Abroug H, Ben Fredj M et al. Management of acute coronary syndrome in emergency departments: a cross sectional multicenter study (Tunisia). *BMC Emerg Med* 2018;**18**:1–9.
36. Hall M, Dondo TB, Yan AT, Goodman SG, Bueno H, Chew DP et al. Association of clinical factors and therapeutic strategies with improvements in survival following non—ST-elevation myocardial infarction, 2003–2013. *JAMA* 2016;**316**:1073–1082.
37. Anderson L, Thompson DR, Oldridge N, Zwisler AD, Rees K, Martin N et al. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2016.
38. Turk-Adawi K, Sarrafzadegan N, Grace SL. Global availability of cardiac rehabilitation. *Nat Rev Cardiol* 2014;**11**:586–596.
39. Grace SL, Turk-Adawi K, Contractor A, Atrey A, Campbell N, Derman W et al. Cardiac rehabilitation delivery model for low-resource settings. *Heart* 2016;**102**:1449–1455.
40. Widimsky P, Fajadet J, Danchin N, Wijns W. Stent 4 Life” targeting PCI at all who will benefit the most. A joint project between EAPCI, Euro-PCR, EUCOMED and the ESC Working Group on Acute Cardiac Care. *EuroIntervention* 2009;**4**:555–557.
41. Ferreira JP, Rossignol P, Dewan P, Lamiral Z, White WB, Pitt B et al. Income level and inequality as complement to geographical differences in cardiovascular trials. *Am Heart J* 2019;**218**:66–74.
42. Dewan P, Rørth R, Jhund PS, Ferreira JP, Zannad F, Shen L et al. Income inequality and outcomes in heart failure: a global between-country analysis. *JACC: Heart Fail* 2019;**7**:336–346.
43. Kaifosova Z, Kala P, Alexander T, Zhang Y, Huo Y, Snyders A et al. Stent for Life Initiative: leading example in building STEMI systems of care in emerging countries. *EuroIntervention* 2014;**10**:T87–T95.
44. European Society of Cardiology. *Prevention in Your Country*. [https://www.escardio.org/Sub-specialty-communities/European-Association-of-Preventive-Cardiology-\(EAPC\)/Advocacy/Prevention-in-your-country](https://www.escardio.org/Sub-specialty-communities/European-Association-of-Preventive-Cardiology-(EAPC)/Advocacy/Prevention-in-your-country).
45. World Health Organization. *Global action plan for the prevention and control of NCDs 2013–2020*. <https://www.who.int/publications/i/item/9789241506236>.